



SCIENCE DIRECT

Register or Login: Password: 
[Home](#) [Search](#) [Journals](#) [Books](#) [Abstract Databases](#) [My Profile](#) [Alerts](#) [? Help](#)
Quick Search:  within [All Full-text Sources](#)  [? Search Tips](#)[← results list](#) [← previous](#) **38 of 40** [next →](#)**Arzneimittel-Forschung**

Volume 33, Issue 5, 1983, Pages 750-754

ISSN: 0004-4172

MEDLINE®

**Tolerability and efficacy of bifonazole in dermatomycoses**

Stettendorf, S

**This Document**[▶ Abstract-MEDLINE](#)**Actions**

- [Cited By](#)
- [Save as Citation Alert](#)
- [E-mail Article](#)
- [Export Citation](#)

**Abstract**

1-[(4-Biphenyl)-phenylmethyl]-1-H-imidazole (bifonazole, Bay h 4502, **Mycospor**)<sup>▶</sup> is a broad spectrum antifungal agent for the topical treatment of dermatomycoses. In addition to tolerability studies, a report is given on the results of 43 clinical studies of efficacy and tolerability involving 1129 patients in various countries. These were placebo-controlled double-blind studies, randomised comparative studies with reference preparations and open studies. They were carried out with bifonazole cream 1%, solution 1%, gel 1% and powder 1%. 58.6% of the patients tested suffered from dermatophytoses, 21.9% cutaneous candidoses, 14.3% from pityriasis versicolor and 5% from other skin infections. The therapeutic efficacy of bifonazole, assessed according to mycological and clinical findings proved good for the indications studied and administered once daily for a treatment period of 2 or 3 weeks. Tolerability - local and systemic - was good. There was no evidence of substance-related side effects. [Clinical Trial, Journal Article, Randomized Controlled Trial; In English; Germany, West]

**CAS Registry Numbers:** Antifungal Agents; Imidazoles; 60628-96-8, bifonazole**Citation Subset Indicators:** Index Medicus

**MeSH Terms:** Adolescent; Adult; Antifungal Agents, adverse effects (AE), \* therapeutic use (TU); Dermatomycoses, \* drug therapy (DT), enzymology (EN), microbiology (MI); Erythrocyte Count; Female; Hemoglobinometry; Humans; Imidazoles, adverse effects (AE), \* therapeutic use (TU); Leukocyte Count; Male; Middle Aged

**Arzneimittel-Forschung**

Volume 33, Issue 5, 1983, Pages 750-754

ISSN: 0004-4172

**This Document**[▶ Abstract-MEDLINE](#)

#### Actions

- [Cited By](#)
- [Save as Citation Alert](#)
- [E-mail Article](#)
- [Export Citation](#)

◀ results list

◀ previous

38 of 40

next ▶

[Home](#)

[Search](#)

[Journals](#)

[Books](#)

[Abstract Databases](#)

[My Profile](#)

[Alerts](#)

[? Help](#)

[Feedback](#) | [Terms & Conditions](#) | [Privacy Policy](#)

Copyright © 2005 [Elsevier B.V.](#) All rights reserved. ScienceDirect® is a registered trademark of Elsevier B.V.